



Pancreas

The pancreas is a gland located in the upper abdomen behind the stomach. Its primary functions are to produce pancreatic juices as an aid to digestion and to manufacture insulin needed for the proper utilization of carbohydrates. It may be the site of an acute inflammatory condition (pancreatitis), abscess, hemorrhage, stones, cyst, pseudocyst, tumor, and cancer.

Abscess and Stone(s)

Present or history, asymptomatic, without complications, fully recovered	Post pone 1 yr. from diagnosis or from any surgical procedure such as debridement, endoscopic retrocyrate cholangio pancreatography (ERCP), or stent placement; then 0.
Others	Individual Consideration

Cyst, Pseudocyst

Cyst or pseudocyst, single, asymptomatic, malignancy ruled out	
Present	Table B
History, resolved, no complications	0
Others: without work-up to exclude malignancy; symptomatic; recurrent or multiple; or with complications such as diabetes, malabsorption, pancreatic insufficiency, etc.	Individual Consideration

This material is intended for insurance informational purposes only and is not personal medical advice for clients. This marketing material includes an expiration date and use of this material must be discontinued as of the expiration date.

FOR INTERNAL USE ONLY. NOT FOR USE WITH THE PUBLIC.

Your Success Matters.

{Name
Phone Number
E-mail Address
Website Address}



Prudential

©2007 The Prudential Insurance Company of America
751 Broad Street, Newark, NJ 07102-3777
Rx099 IFS-A060831 Ed. 06/07 Exp. 12/08

Pancreatitis

Pancreatitis may be acute or chronic and attacks are usually manifested by severe abdominal pain. While stones in the bile duct may cause acute pancreatitis, the most common cause of chronic pancreatitis is alcoholism.

Not due to alcohol or malignancy; no surgical procedures such as drainage per endoscope (ERCP) or pancreatectomy and no complications such as diabetes, pseudocyst, or malabsorption	
Acute pancreatitis, recovered and asymptomatic within 8 weeks, 1-2 episodes only, cause identified and corrected	0
Chronic (lasting more than 8 weeks) or recurrent (more than two acute episodes) pancreatitis	
Within 6 months of last occurrence	Post pone
6-12 months	Table G
2nd year	Table E
3rd year	Table C
4th year	Table B
Thereafter	0
With surgical treatment such as pancreatectomy or endoscopic drainage, now recovered	Rate as chronic pancreatitis
Others: with complications, no work-up to rule out malignancy	Decline
Due to alcohol, now abstinent and without complications such as diabetes, pseudocyst, malabsorption, etc.	
1-2 episodes of acute pancreatitis, recovered and asymptomatic	Rate for alcohol abuse See Rx for Success #102
For more than two episodes or for chronic symptoms, applicant would be rated as below or rated for alcohol abuse if the latter is higher.	
Within 2 years of diagnosis	Post pone
3rd - 5th years	Table G
6th - 10th years	Table C
Thereafter	Table B
Others	Decline

This material is intended for insurance informational purposes only and is not personal medical advice for clients.

This marketing material includes an expiration date and use of this material must be discontinued as of the expiration date.

FOR INTERNAL USE ONLY. NOT FOR USE WITH THE PUBLIC.

